

Operators which factor through Banach lattices not containing c_0

by

N. Ghoussoub and W. B. Johnson*

In this supplement to [GJ1], [GJ3], we give an intrinsic characterization of (bounded, linear) operators on Banach lattices which factor through Banach lattices not containing a copy of c_0 which complements the characterization of [GJ1], [GJ3] that an operator admits such a factorization if and only if it can be written as the product of two operators neither of which preserves a copy of c_0 . The intrinsic characterization is that the restriction of the second adjoint of the operator to the ideal generated by the lattice in its bidual does not preserve a copy of c_0 . This property of an operator was introduced by C. Niculescu [N2] under the name “strong type B”. That this condition is stronger than the property that the operator itself does not preserve a copy of c_0 is exhibited by the example constructed in [FGJ] and studied further in [GJ2].

When we wrote [GJ1], [GJ3] we did not know that Niculescu [N1], [N2], [N3] was independently considering some of the same problems; we now want to acknowledge his priority for discovering some results that overlap with ours. Also, in remark 1 below, we indicate how our results, when translated into Niculescu’s language, answer some questions he raised.

Before stating the theorem of this note, we introduce some notation: Given a Banach lattice E , $I(E)$ (respectively, $Bd(E)$) denotes the closed order ideal (respectively, band) in E^{**} generated by E . The symbols E, F, G will be reserved for Banach lattices while X, Y, Z will be used for general Banach spaces. Terminology is as in [GJ3], and the reader will probably need a copy of [GJ3] on hand to follow the proof of the theorem.

Theorem. *Let E be a Banach lattice and let T be an operator from E into a Banach space X . The following are equivalent:*

- (1) $T^{**}|_{I(E)}$ does not preserve a copy of c_0 .
- (2) $T^{**}|_{I(E)}$ does not preserve a positive disjoint copy of c_0 .
- (3) $T^{**}[Bd(E)] \subset X$.
- (4) There exists a Banach space Y and bounded linear operators $u: E \rightarrow Y$ and $v: Y \rightarrow X$ such that neither u nor v preserves a copy of c_0 and $T = vu$.
- (5) There exists a weakly sequentially complete Banach lattice F , an interval preserving lattice homomorphism $R: I(E) \rightarrow F$, and an operator $S: F \rightarrow X$ so that $T^{**}|_{I(E)} = SR$.

Proof. That (1) and (2) are equivalent follows from Theorem I.3 of [GJ3] (that the other conditions in Theorem I.3 are equivalent to (2) was proved earlier by Niculescu—see Theorem 3.4 in [N2]). Niculescu [N2, Proposition 4.2] stated the equivalence of (2) and (3); we follow his lead in leaving the proof as an exercise for the reader, but note that (2) is the statement we tie to the other conditions. The equivalence of (4) with (5) was proved in Corollary I.8 in [GJ3] and (5) $\Rightarrow$ (2) is immediate, so we only need to check that (2) implies (5). In fact, the proof of this implication is essentially contained in the proof

* Supported in part by NSF DMS-8703815

of Theorem I.7 of [GJ3], but since only weaker implications are explicitly stated there we present here a detailed proof.

Let W be the absolute convex solid hull of $T^*Ball(X^*)$ in E^* and let G be the interpolation space (cf. [GJ3]) corresponding to W in E^* . Let $J: G \rightarrow E^*$ be the canonical injection and $I: X^* \rightarrow G$ such that $T^* = JI$. By Lemma I.5 of [GJ3] we have for each μ in E^{**} and $n = 1, 2, \dots$,

$$\|J^*\mu\| \leq 2^n \sup \{ \|T^{**}\nu\|: \nu \in E^{**}, |\nu| \leq |\mu| \} + 2^{-n}\|\mu\|. \quad (\dagger)$$

Letting F be the norm closure of $J^*[I(E)]$ in G^* and setting $R = J^*|_{I(E)}$, $S = I^*|_F$, we have that $T^{**}|_{I(E)}$ factors through R and hence in order to complete the proof we only need to check that F does not contain a copy of c_0 .

Since $I(E)$ is an ideal in E^{**} we have from $(\dagger)$ that for each $f \in I(E)$ and $n = 1, 2, \dots$,

$$\|Rf\| \leq 2^n \sup \{ \|T^{**}g\|: g \in I(E), |g| \leq |f| \} + 2^{-n}\|f\|. \quad (\ddagger)$$

Now J^* is a lattice homomorphism since J is interval preserving ([AB, p. 90]). Since $T^{**}|_{I(E)}$ does not preserve a copy of c_0 , neither does R by Lemma I.1.a of [GJ3], hence $(\ddagger)$ implies that F has order continuous norm by Lemma I.1.b of [GJ3].

We first prove that if $(e_m)_{m=1}^\infty$ is an increasing sequence in $I(E)_+$ such that $(J^*e_m) = (Re_m)$ is norm bounded in F then it is norm convergent in F . Indeed, let μ be the w^* -limit of $(J^*e_m)_{m=1}^\infty$ in G^* . Since $J^*[E^{**}]$ is norm dense in G^* , there exists for each $\epsilon > 0$ an element $\nu_\epsilon \in E^{**}$ such that $\|J^*\nu_\epsilon - \mu\| \leq \epsilon$.

Since $I(E)$ is an ideal in E^{**} , the sequence $(\nu_\epsilon \wedge e_m)_{m=1}^\infty$ is in $I(E)$; moreover, it is increasing and norm bounded by $\|\nu_\epsilon\|$, hence $J^*(\nu_\epsilon \wedge e_m)$ norm converges to some z_ϵ in F by Theorem I.3.d in [GJ3]. Using again the fact that J^* is a lattice homomorphism, we have that $J^*(\nu_\epsilon \wedge e_m) = J^*(\nu_\epsilon) \wedge J^*(e_m)$ hence $J^*(\nu_\epsilon \wedge e_m)$ w^* -converges in G^* to $J^*(\nu_\epsilon) \wedge \mu$. Therefore $J^*(\nu_\epsilon) \wedge \mu = z_\epsilon$ is in F . Thus we get

$$\lim_{\epsilon \rightarrow 0} \|z_\epsilon - \mu\| = \lim_{\epsilon \rightarrow 0} \|(J^*\nu_\epsilon) \wedge \mu - \mu\| \leq \lim_{\epsilon \rightarrow 0} \|J^*\nu_\epsilon - \mu\| = 0.$$

It follows that μ is in F . Since the sequence $(J^*e_m)_{m=1}^\infty$ is in the order interval $[0, \mu]$ and F is order continuous, it follows that $(J^*e_m)_{m=1}^\infty$ norm converges to μ .

Suppose now that $(f_m)_{m=1}^\infty$ is any increasing sequence in F . To show that it is norm convergent, it is enough to show that for any $\epsilon > 0$ there exists $(e_m)_{m=1}^\infty$ increasing in $I(E)_+$ such that $\|f_m - J^*e_m\| \leq \epsilon$. For that, let $z_m \in I(E)_+$ be such that $\|J^*z_m - f_m\| \leq \epsilon 2^{-m}$. Set $e_m = \vee_{i=1}^m z_i$. Then $(e_m)_{m=1}^\infty$ is increasing and

$$|J^*e_m - f_m| = |J^*(\vee_{i=1}^m z_i) - f_m| = |\vee_{i=1}^m J^*z_i - \vee_{i=1}^m f_i| \leq \sum_{i=1}^m |J^*z_i - f_i|.$$

It follows that

$$\|J^*e_m - f_m\| \leq \sum_{i=1}^m \|f_i - J^*z_i\| \leq \epsilon \sum_{i=1}^m 2^{-i} \leq \epsilon.$$

By the classical version of Theorem I.3 in [GJ3], F does not contain a copy of c_0 and hence F is a weakly sequentially complete Banach lattice ([LT, p. 34]). ■

In the above proof of (5) $\Rightarrow$ (2), if X is Banach lattice and T is a positive operator, then every positive operator dominated by T also factors through R . Thus we get:

Corollary. *Let E be a Banach lattice, let T be a positive operator from E into a Banach lattice X , and assume that $T^{**}|_{I(E)}$ does not preserve a positive disjoint copy of c_0 . Then there exists a weakly sequentially complete Banach lattice F and an interval preserving lattice homomorphism $R: I(E) \rightarrow F$ such that for any operator $0 \leq T_1 \leq T$ there is a positive operator $S_1: F \rightarrow X$ so that $T_1^{**}|_{I(E)} = S_1 R$.*

Remark 1. The examples in [FGJ] of positive quotient mappings onto c_0 which do not preserve a copy of c_0 were shown in [GJ2] to fail property (4) of the theorem, hence are, in Niculescu's terminology [N2], operators of type B which are not of strong type B. This answers problem 4.1 in [N2]. (However, if E is order continuous then $I(E) = E$ and so operators from E of type B are of strong type B.) These operators also give a solution to problem 4.4 in [N2]; namely, there exists a lattice homomorphism T on some Banach lattice which does not preserve a copy of c_0 or of l_1 , yet T^2 is not weakly compact, while T^3 must be weakly compact: see remark 2 in [GJ3, p. 164].

Remark 2. One of the operators mentioned in remark 1 (" T_1 " in the terminology of [GJ2]) maps weakly Cauchy sequences into weakly convergent sequences; this gives a counterexample to one part of Proposition 4.2 in [N2].

Remark 3. Since a weakly sequentially complete Banach lattice has the so-called *analytic Radon- Nikodym—ARN—property* (cf. [BD]), an operator which satisfies condition (3) of the theorem is an ARN-operator. This implication was recently proved in a direct way by Bukhvalov [B]. Although aware of [GJ3], Bukhvalov mistakenly thought that his result could not follow from the factorizations in [GJ3] (he applied the error from [N2] mentioned in remark 2 to conclude that the operators mentioned in remark 1 satisfy condition (3) of the theorem).

References

- [AB] C. D. Aliprantis and O. Burkinshaw, *Positive operators*, **Academic Press** (1984).
- [BD] A. V. Bukhvalov and A. A. Danilevich, *Boundary properties of analytic and harmonic functions with values in Banach space*, **Mat. Zametki** **31** (1982), 203–214. English translation: **Math. Notes** **31** (1982), 104–110.
- [B] A. V. Bukhvalov, *On the analytic Radon- Nikodym property*, [unpublished manuscript].
- [FGJ] T. Figiel, N. Ghoussoub, and W. B. Johnson, *On the structure of non-weakly compact operators on Banach lattices*, **Math. Ann.** **257** (1981), 317–334.
- [GJ1] N. Ghoussoub and W. B. Johnson, *On subspaces of Banach lattices not containing $C(\Delta)$* , [unpublished].
- [GJ2] N. Ghoussoub and W. B. Johnson, *Counterexamples to several problems on the factorization of bounded linear operators*, **Proc. AMS** **92** (1984), 233–238.

- [GJ3] N. Ghoussoub and W. B. Johnson, *Factoring operators through Banach lattices not containing $C(0, 1)$* , **Math. Zeit.** **194** (1987), 153–171.
- [LT] J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces II: Function spaces*, **Ergebnisse Math. Grenzgebiete** **97** Springer-Verlag (1979).
- [N1] C. Niculescu, *Weak compactness in Banach lattices*, **J. Operator Theory** **6** (1981), 217–231.
- [N2] C. Niculescu, *Order σ -continuous operators on Banach lattices*, **Lecture Notes in Math.** **991** (1983), 188–201.
- [N3] C. Niculescu, *Operators of Type A and local absolute continuity*, **J. Operator Theory** **13** (1985), 49– 61.

Department of Mathematics, University of British Columbia, Vancouver, B. C., Canada

Department of Mathematics, Texas A&M University, College Station, TX 77843, USA